

DETERMINING THE CRITICAL NEW SNOW DEPTH FOR A DESTRUCTIVE AVALANCHE BY CONSIDERING THE RETURN PERIOD

Jürg Schweizer*, Christoph Mitterer and Lukas Stoffel
WSL Institute for Snow and Avalanche Research SLF, Davos, Switzerland

ABSTRACT: Snow avalanche danger assessment for avalanche paths threatening a highway or a residential area is usually related to new snow depth. Given the extensive protection work in the Alps, the avalanche control service (also called avalanche commission) responsible for danger assessment will usually monitor the avalanche situation throughout the winter, but only become active in case of a major snow fall. Related safety concepts describing the procedures and measures to be taken in a given danger situation are therefore typically based on threshold values for new snow. By analysing the avalanche occurrence of a major avalanche path, we show that the return period of an avalanche to, for example, the road is about 5 years, whereas the return period for the corresponding new snow depth is substantially smaller, in our case slightly less than 2 years. Similar proportions were found for a number of other avalanche paths with different snow climate. The return period of the critical new snow depth is typically about 2-5 times smaller than the return period of the avalanche. This proportion is expected to increase with increasing return period. Hence, based on the return period of an avalanche path a first estimate for the critical new snow depth can be made. With a return period of the critical new snow depth of 1-2 years, avalanche prediction for individual avalanche path becomes very challenging since the false alarm ratio is expected to be high.

KEYWORDS: snow avalanche, snow stability evaluation, avalanche forecasting

1. INTRODUCTION

In the European Alps most of the severe avalanche problems have been mitigated in the past decades by permanent protection measures such as supporting structures in the starting zone, or dams and sheds in the run-out zone. Still, there remain very many avalanche paths without permanent protection measures in place. They either produce infrequent events or permanent protection works is technically difficult to implement and/or too costly (poor cost-to-benefit ratio). In particular for economic reasons, avalanche forecasting (i.e. preventive closures) – often combined with artificial avalanche release is now frequently favoured over permanent protection works.

This solution requires a well organized local avalanche control service with personnel who is usually hired part time and very often are well qualified volunteers. As critical situations are infrequent, the avalanche service has to assess the situation and take action only occasionally during

the winter. Although it is recommended that they closely follow the avalanche situation during the winter, it is common that a service only starts working when a major snow storm is pre-announced (Stoffel and Schweizer, 2008). Ideally, the avalanche service has established a concept that connects a given avalanche situation to some temporary protection measures. The avalanche situation in such a safety concept is often characterized by the amount of snow loading. Threshold values are commonly determined based on past events. Often the non-events are not considered in this type of analysis. These critical values should be considered as a first guess and always be adapted to the actual situation. After an unexpected event, the snow loading is often considered as relatively minor and not comparable to the often large extent of the unexpected destructive avalanche.

Considering the return period for hazard mapping is common (e.g. Ancey et al., 2004; Burkard and Salm, 1992), but the approach is rarely combined with local avalanche prediction, except in the pioneering work by Föhn and Meister (1982). Avalanche forecasting tools for one or more individual avalanche path do not exist, probably due to the fact that rare events do not allow developing a statistical forecasting model. Avalanche activity around Zuoz (Engadine valley, Switzerland) has been related to snow and snow-pack parameters (Stoffel et al., 1998). They found

Corresponding author address: Jürg Schweizer,
WSL Institute for Snow and Avalanche Research SLF, Flüelastrasse 11,
CH-7260 Davos Dorf, Switzerland;
tel: +41 81 4170164; fax: +41 81 4170110;
email: schweizer@slf.ch

that new snow depth alone was insufficient for forecasting, but that snowpack stratigraphy and temperature evolution were essential contributing factors even for large catastrophic avalanches. A similar analysis for various regions was performed by Schneebeli et al. (1998).

One of the snow loading variables, the increase in snow depth during 3 days ($\Delta HS3d$) is also used in the context of modelling the dynamics of large catastrophic avalanches for hazard mapping to assess the fracture depth. This approach has been questioned (e.g. Barbolini et al., 2002).

The aim of this study is to analyse the avalanche activity in an active avalanche path (Salezertobel, Davos, Switzerland), derive values for the critical new snow depth and relate the return period of the critical new snow depth to the avalanche return period. We will then describe the ratio of return periods for a number of other avalanche path in order to provide some rough guidance on how to establish preliminary threshold values for an avalanche path where little information is available apart from avalanche occurrence data.

2. METHODS AND DATA

The analysis was made for an avalanche path that runs towards the main road that enters the city of Davos (1560 m a.s.l., Eastern Swiss Alps) from the north: the Salezertobel path. The Salezertobel avalanche path has already been analysed by Föhn and Meister (1982). The starting zone reaches up to 2500 m a.s.l., is about 33-37° steep and has mainly easterly to south-easterly aspect. The distance to the road is about 1800 m. Avalanche records go back to the 15th century. For the last about 60 years the occurrence was consistently recorded. However, the avalanche extent is not always known and there were many small events. We will consider the winter periods from 1950-1951 to 2007-2008 (58 years). About 70 avalanches were recorded. We will only consider the 55 avalanche events that were more or less well documented. Except for one event, the avalanches were mapped and available for GIS analysis. From the 55 avalanches considered, 34 were large events that had a runout below 1700 m a.s.l., i.e. on the alluvial fan above the road. From these large events 12 reached (± 20 m) the road or the shed (since 1984 the road is protected by a snow shed; construction started in 1981). In five winters two (and once even three) large events were recorded, still we consider all large avalanches as independent events.

The snow and weather data used for the analysis were recorded at the study plots of Weissfluhjoch (2540 m a.s.l.) above Davos and of Davos Dorf (1560/1590 m a.s.l.). At both locations new snow depth was recorded daily on a snow board. Other meteorological parameters included air temperature, wind speed, radiation etc. For the analysis, we used daily values of the 58 winters from 1 November to 30 April, in total 10,513 daily records. To simplify the analysis we reduced the dataset and only considered days with a new snow depth $HN \geq 10$ cm measured at Weissfluhjoch ($N = 1540$). Furthermore, for days immediately after an event, the 3, 5 or 10 day sum of new snow depth was not considered.

Snow stratigraphy was included based on the bi-weekly snow profiles taken at the Weissfluhjoch, Büschalp (1960 m a.s.l.) and Davos Dorf study plots. Profiles were classified according to Schweizer and Wiesinger (2001) into those with weak basal layers (profile types 1-5) and those with well consolidated (rather hard) basal layers (profile types 6-10). This snow stratigraphy classification was only available for analysis for days with large avalanche events.

To contrast variables from days with no avalanche events to avalanche days the non-parametric Mann-Whitney *U*-Test was used. Categorical data such as snowpack classification were cross-tabulated and a Yates' corrected Pearson χ^2 statistic was calculated. A level of significance $p = 0.05$ was chosen to decide whether the observed differences were statistically significant. Split (or threshold) values between two categories were determined with the classification tree method (Breiman et al., 1998).

To characterise the return period of a given new snow amount (e.g. new snow depth of 24 hours: HN , or 3-d sum of new snow depth: $HN3d$), we used the Gumbel extreme value statistics.

3. RESULTS

Figure 1 shows the avalanche frequency in the Salezertobel path for the 54 avalanches mapped from 1950-1951 to 2007-2008. During the 58 years considered, 12 avalanches reached the road (± 20 m) so that the return period for an avalanche to the road is about 5 years.

We will first consider the meteorological situation for the 12 avalanches that reached the road. The new snow amount prior to the release varied widely. For example, the 3-d sum of new snow depth at Weissfluhjoch varied between 51 cm and 127 cm, with a median value of 68 cm.




Figure 1: Avalanche frequency in the Salezertobel path (Davos, Switzerland). During a period of 58 years (1950-1951 to 2007-2008) 54 avalanche events were mapped.

For nine out of 12 avalanches $HN3d$ measured at Weissfluhjoch was at least 64 cm (1st quartile). Considering the new snow measurements at the valley bottom showed that the median $HN3d$ was 67 cm with a range of 22 cm to 110 cm. For nine out of 12 avalanches $HN3d$ measured at Davos was at least 41 cm. The air temperature change to the previous day ΔT_a was in most cases positive at Weissfluhjoch, i.e. about +3°C. There was no comparable trend for the temperature measured at Davos. Analysing the snow stratigraphy showed that the profiles taken prior to the release at the lower elevation study plots (Davos and Büschalp) had almost exclusively a weak basal snowpack layers; at the elevation of the Weissfluhjoch eight out of 12 profiles had a weak base.

In the following, we compare the above described conditions for the avalanches that hit the road to those of the other 22 rather large events. All variables related to snow loading (HN , $HN3d$, $HN5d$, $\Delta HS3d$) showed larger median values for the avalanches that hit the road compared to the other large events (either measured at Weissfluhjoch or at Davos) (Figure 2); the differences were statistically significant (Mann-Whitney U -test; $p \leq 0.015$) except for $HN5d$ measured at Davos ($p = 0.052$). The level of significance p was in general lower for the values measured at Davos. Based on univariate tree statistics, a




Figure 2: Sum of new snow depth for 3 and 5 days ($HN3d$, $HN5d$), and 3-day increase in snow depth ($\Delta HS3d$) measured at (a) the Weissfluhjoch and (b) Davos for large avalanches that stopped above the road (left, in blue; $N = 22$) and avalanches that hit the road (right, in red; $N = 12$).

threshold value for an avalanche to the road of ≥ 57 cm and ≥ 59 cm for $HN3d$ was found, measured at Weissfluhjoch and Davos, respectively. Also significant variables were ΔT_a at Weissfluhjoch ($\geq 2.1^\circ\text{C}$, $p = 0.022$) and the snow depth 3 days before the event ($HS3d$) at Weissfluhjoch (≥ 140 cm, $p = 0.035$). The proportion of profiles (taken at either Büschalp or Davos) with weak base was significantly larger for days with avalanches that hit the road. Whereas a strong base suggests that an avalanche will not reach the road, a weak base has no discriminating power. Though, for almost all avalanches that reached the road a snowpack with a weak base existed – but weak snowpack basal layers existed also when many of the large avalanches stopped above the road.

Based on the analysis of the events only, a new snow amount of about 55-60 cm (measured either on the level of the starting zone at the Weissfluhjoch or in the valley bottom at Davos) seems to indicate that an avalanche might reach the road. The return period of a new snow depth of about 55-60 cm in 3 days at Davos is about 1.5-2 years, at Weissfluhjoch it is about 1 year.

Table 1: Optimal threshold values based on unweighted average accuracy to discriminate between days when a large avalanche occurred and non-event days.

Loading variable	Threshold value	unweighted average accuracy
$HN3d_{WFJ}$	≥ 45 cm	71.3%
$HN5d_{WFJ}$	≥ 54 cm	69.1%
$HN3d_{DAV}$	≥ 33 cm	72.5%
$HN5d_{DAV}$	≥ 38 cm	72.7%




Figure 3: Sum of new snow depth for 3 days measured at (a) the Weissfluhjoch and (b) Davos for all days (left, in blue; $N = 1528$) and avalanches that hit the road (right, in red; $N = 12$).

Next, we will consider if it is possible to forecast whether a large avalanche has to be expected. We compared the snow and weather situation at days ($N = 34$) when a large avalanche occurred to those days when there was no avalanche. All loading parameters were significant variables ($p \leq 0.01$) except for *HN* at Weissfluhjoch ($p = 0.25$). Also significant variables were the

snow depth variables (*HS*, *HS3d*). Tree statistics with standard parameters did not suggest any split value for any of the snow loading parameters, except for *HN* at Davos (≥ 49 cm). However, as the dataset was very unbalanced it is not surprising that no split values were found, and the one found is questionable. Hence, we optimized the thresholds to reach the highest unweighted average accuracy or the highest true skill score. Results are summarized in Table 1. The unweighted average accuracy can be increased to 75% if in addition to *HN3d* or *HN5d* at Davos also the snow depth *HS* (≥ 76 cm) is considered. However, the probability of detection (correct avalanche events) is only about 65%, whereas the false alarm ratio is about 90%. In other words, the scheme is not applicable in practice.

Forecasting the avalanches that reached the road was slightly easier, since the snow and weather situation prior to the release was in general more extreme (Figure 3). We again primarily considered the loading variables (Table 2). First of all, the results show that the false alarm ratio (FAR) was generally higher than 90%, i.e. only in 1 (or even less) out of 10, for example, road clo-

Table 2: Classification models to discriminate between days when a large avalanche occurred and non-event days with corresponding skill scores.

Variable	Threshold value	Probability of detection (POD)	True skill score (HK)	Unweighted average accuracy	False alarm ratio (FAR)
<i>Weissfluhjoch</i>					
<i>HN</i>	≥ 33 cm	66.7%	55.7%	77.9%	95.5%
<i>HN3d</i>	≥ 51 cm	100%	78.3%	89.2%	96.6%
<i>HN5d</i>	≥ 59 cm	100%	75.3%	87.7%	97.0%
<i>Davos</i>					
<i>HN</i>	≥ 36 cm	58.3%	54.9%	77.3%	89.1%
<i>HN3d</i>	≥ 42 cm	75.0%	63.0%	81.5%	95.4%
<i>HN5d</i>	≥ 44 cm	83.3%	65.5%	82.7%	96.5%
<i>Combinations</i>					
<i>HN3d_WFJ</i> and <i>HS_DAV</i>	≥ 59 cm ≥ 76 cm	83.3%	74.6%	87.3%	93.2%
<i>HN3d_WFJ</i> and <i>HS_DAV</i>	≥ 64 cm ≥ 76 cm	75.0%	68.0%	84.0%	92.4%
<i>HN3d_DAV</i> and <i>HS_DAV</i>	≥ 42 cm ≥ 76 cm	75.0%	66.6%	83.3%	93.6%
<i>HN3d_DAV</i> and <i>HS_DAV</i>	≥ 59 cm ≥ 76 cm	66.7%	62.0%	81.0%	90.1%
<i>HN3d_DAV</i> and <i>HS_DAV</i> and <i>HN_DAV</i>	≥ 59 cm ≥ 76 cm ≥ 33 cm	66.7%	64.8%	82.4%	78.4%

sure an avalanche actually released. Obviously, the false alarm ratio decreased with increasing threshold value (or increasing number of variables), but inevitably more avalanches were missed, i.e. the number of “hits”, or the probability of detection (POD) decreased. If we would, for example, request that a classification model has to have a $POD \geq 75\%$ and a $FAR < 90\%$, none of the models in Table 2 would pass the test. The variables at Davos had slightly more discriminating power as the new snow amounts for the avalanche events hitting the road were more extreme in Davos than at the Weissfluhjoch. In addition, the false alarm ratio was slightly lower if the loading variables measured in the valley bottom were used for similar threshold values.

In summary, the analysis suggests that a critical new snow depth ($HN3d_{crit}$) of about 55-60 cm if measured at Davos and about 60 cm if measured at Weissfluhjoch seems appropriate. However, these values imply that about every third avalanche to the road would be missed. Still, most of the times when the road would be closed, no avalanche would release. In fact, a critical new snow depth of 60 cm has a return period at Weissfluhjoch of slightly more than 1 year (Gumbel statistics), but occurs about four times per winter. Given this threshold, the road should be closed (if there would be no shed) at least four times per winter, but an avalanche would reach the road only every 5 years. Due to the high false alarm rate forecasting based on the threshold value of 60 cm from Weissfluhjoch seems not feasible. Using the snowfall data measured at Davos is somewhat more appropriate. A critical new snow

depth of 55-60 cm only occurs slightly less than about twice per winter (1.8 times per winter). In about 30% of these cases the snow depth is less than 80-100 cm at the elevation of the valley bottom so that an avalanche hitting the road is unlikely. Hence, a critical situation is reached only about 1.3 times per winter.

A similar, though less detailed analysis was performed for other sites where an avalanche path threatens a road or communication line. In these cases the occurrence record is less complete and often only the major events that reached the road were recorded. Consequently, the threshold values cannot be determined statistically by comparing events to the road to events that stopped above the road, nor by comparing conditions of the avalanche events with those of the non-events. The critical new snow depth as measured in a study plot in the valley bottom (usually within a couple kilometres from the starting zone) corresponds to about the 10-30% percentile depending on the observation period and the number of recorded events. Table 3 compiles these results. The snow depth in the valley bottom was usually >50-60 cm at the beginning of the snowfall period.

The return period of the critical new snow depth was about 2-5 times smaller than the return period of the avalanche event. It is expected that with increasing return period the ratio might increase, i.e. even for the very rare and extreme events the critical new snow depth will often not be extraordinary. This possible trend is reflected in the few examples shown in Table 3. For avalanche path with a return period of about 5 years

Table 3: Return periods of avalanche events that threaten a communication line and of corresponding critical new snow depth ($HN3d_{crit}$).

Site	Avalanche return period (years)	Critical new snow depth $HN3d_{crit}$ (cm)	Return period $HN3d_{crit}$ (=potential damage) (years)	Return period ratio
Salezertobel (Davos)	5	55-60	1.5-2	~3
Breitzug (Davos)	5	65	2	2.5
Gonda, Lavin (Eastern Swiss Alps)	5	65	2.5	2
Zuoz (Engadine)	5	35-40	1-2	~3
Col du Pillon (Les Diablerets, Western Swiss Alps)	10	70-80	~2	~5
Ravaisch (Samnaun, Eastern Swiss Alps)	12	70	2.5	~5
Kreuzbachtobel (Pfäfers-Vättis, Northern Swiss Alps)	20	80	4	5

the ratio was about 2-3, whereas for the path with a return period of 10-20 years, the ratio was higher, that is about 5. Based on the limited data set analysed no relation to the snow climate can be found.

In Table 3 only single avalanche path were considered. If several avalanche path with similar return period endanger a road the combined return period is lower than the return period in the individual path whereas the return period of the critical new snow depth will be the same so that the ratio will be lower, probably about 1-2.

4. DISCUSSION

If we assume that the return period of the critical new snow depth is about 2-5 times smaller than the return period of the avalanche event under consideration – as our preliminary analysis suggests, we can estimate the critical new snow depth from nearby snow observations in the same area having the same snow climate even if there is no information about the snow and weather conditions at the times of the avalanche events. For example, for some avalanche path only the year might be none when an avalanche hit the road.

Given this information we can estimate the frequency the road is threatened by a potential avalanche release. We expect this estimate to usually be more specific and hence useful (about ± 10 cm within the observed critical value) than what is indicated in rough guidelines on the relation between new snow depth and avalanche activity (e.g. Salm, 1982). A typical range in these guidelines for the problems listed in the Table 3 is 50-80 cm of new snow. If the estimate of the critical new snow depth indicates that the road might be threatened many times per winter this implies that reliable forecasting might be impossible – and permanent avalanche protection works might be better suited to solve the avalanche problem under consideration.

Certainly, this proposal is preliminary, but the focus on return periods also shows some of the challenging problems inherent to the forecasting of large avalanches in a given avalanche path.

5. CONCLUSIONS

We have analysed the avalanche activity for the well documented Salezertobel avalanche path near Davos (Switzerland) for the period 1950-1951 to 2007-2008. The return period for an avalanche to the road level (now protected by a shed)

was 5 years. These large avalanche events were all related to substantial snow loading, a snow depth above terrain roughness, a snow stratigraphy which was characterised at the elevation of the track and the run-out zone by weak basal layers and a slightly increasing air temperature trend. However, when including the non-event days in the analysis, forecasting based on the above characteristics becomes difficult due to the high number false alarms. Simple classification models based on $HN3d$ and HS measured at Davos showed that the critical new snow depth for an avalanche to the road level is about 55-60 cm.

This value has a return period of less than two years. For return periods of a few years, the Gumbel statistics largely overestimate the return period. Consequently, there were many days when the critical new snow depth was reached so that the number of false alarms was so high that a model simply based on a critical new snow amount is not applicable in practice. The number of false alarms was reduced by considering one or two additional variables such as the snow depth. Still, the probability that an avalanche reaches the road when the model suggest so, was ≤ 0.15 . Obviously, in operational avalanche forecasting for roads or residential areas, many other variables are considered and the critical new snow depth is adapted so that most avalanche services perform significantly better than our simple models.

For the analysis, we used snow and weather data from two locations: one representative for the starting zone (Weissfluhjoch), the other for the run-out zone (Davos). The data collected at the valley bottom were as useful as the data from the elevation of the starting zone. Forecasting based on data from Weissfluhjoch – though highly correlated with large avalanche events – caused more false alarms than when the data from Davos were used. Though the conditions at the elevation of the starting zone are undoubtedly better captured with automatic stations at this elevation, the data might not be appropriate for forecasting extreme events due to their inherent low predictability.

The difference in snow depth over the last three days $\Delta HS3d$ was a significant variable to forecast an avalanche to the road, but by far not the best one. However, for avalanches that reached the road level $\Delta HS3d$ was significantly larger than for avalanches that had a shorter run-out, indicating that the run-out distance is related to $\Delta HS3d$ – which is commonly assumed in avalanche dynamics calculations.

The ratio of the return period of the critical new snow depth to the avalanche return period

was evaluated for six more avalanche paths and values in the range of 2 to 5 were found. This finding might be useful to preliminarily assess the critical new snow depth for an avalanche path for which only the return period might be known.

Though avalanche control services are in general probably more successful than a simple model based on a critical new snow depth, the generally low predictability makes the prediction of an avalanche event in a specific avalanche highly uncertain. Therefore, avalanche forecasting (i.e. for example, preventive road closures) – even when combined with explosive control might not always be the best option when evaluating the cost effectiveness of potential avalanche protection measures (costs vs. prevented death).

ACKNOWLEDGEMENTS

We would like to thank Roland Meister for sharing his expertise and data on the Salezertobel avalanche path and Stephan Margreth for valuable input.

REFERENCES

- Ancey, C., C. Gervasoni, and M. Meunier, 2004: Computing extreme avalanches. *Cold Reg. Sci. Technol.*, **39**(2-3), 161-180.
- Barbolini, M., L. Natale, and F. Savi, 2002: Effects of release conditions uncertainty on avalanche hazard mapping. *Natural Hazards*, **25**, 225-244.
- Breiman, L., J.H. Friedman, R.A. Olshen, and C.J. Stone, 1998: *Classification and regression trees*. CRC Press, Boca Raton CA, U.S.A., 368 pp.
- Burkard, A. and B. Salm, 1992: Die Bestimmung der mittleren Anrissmächtigkeit d_0 zur Berechnung von Fließlawinen. *Eidg. Institut für Schnee- und Lawinenforschung, Internal Report No. 668*, Weissfluhjoch-Davos, Switzerland, 16 pp.
- Föhn, P. and R. Meister, 1982: Determination of avalanche magnitude and frequency by direct observations and/or with aid of indirect snowcover data. *Proceedings IUFRO/FAO colloquium on research on small torrential watersheds (incl. avalanches)*, Grenoble, France, June 1981, Mitteilungen der forstlichen Bundesversuchsanstalt, Vienna, Austria, Vol. 144, 207-228.
- Salm, B., 1982: *Lawinenkunde für den Praktiker*. Verlag des Schweizer Alpen-Club SAC, Berne, Switzerland, 148 pp.
- Schneebeli, M., M. Laternser, P. Föhn, and W. Ammann, 1998: *Wechselwirkungen zwischen Klima, Lawinen und technischen Massnahmen*. Nationales Forschungsprogramm "Klimaänderungen und Naturkatastrophen" (NFP 31), Schlussbericht, vdf Hochschulverlag an der ETH Zürich, Zürich, Switzerland, 134 pp.
- Schweizer, J. and T. Wiesinger, 2001: Snow profile interpretation for stability evaluation. *Cold Reg. Sci. Technol.*, **33**(2-3), 179-188.
- Stoffel, A., R. Meister, and J. Schweizer, 1998: Spatial characteristics of avalanche activity in an alpine valley - a GIS approach. *Ann. Glaciol.*, **26**, 329-336.
- Stoffel, L. and J. Schweizer, 2008: Guidelines for avalanche control services: organisation, hazard assessment and documentation – an example from Switzerland. *Proceedings ISSW 2008. International Snow Science Workshop, Whistler BC, Canada, 21-26 September 2008*, this issue.